

Adaptive MLMC for Computing Probabilities

Abdul-Lateef Haji-Ali

Jonathan Spence (HWU)

Aretha Teckentrup (University of Edinburgh)

Heriot-Watt University

MCM — August 17, 2021

The problem: Computing probabilities

$$\mathbb{P}[Z \in \Omega] = \mathbb{E}[\mathbb{I}_{Z \in \Omega}]$$

where Z is a d -dimensional random variable and $\Omega \in \mathbb{R}^d$. This problem can be written in the form

$$\mathbb{P}[X > 0] = \mathbb{E}[\mathbb{I}_{X > 0}]$$

for a one-dimensional random variable X which is the signed distance of Z to Ω .

The problem: Computing probabilities

$$\mathbb{P}[Z \in \Omega] = \mathbb{E}[\mathbb{I}_{Z \in \Omega}]$$

where Z is a d -dimensional random variable and $\Omega \in \mathbb{R}^d$. This problem can be written in the form

$$\mathbb{P}[X > 0] = \mathbb{E}[\mathbb{I}_{X > 0}]$$

for a one-dimensional random variable X which is the signed distance of Z to Ω .

Two main reasons this problem can be challenging:

- 1 The event is rare,
- 2 and the complexity of sampling X .

The problem: Computing probabilities

- Financial risk assessment $X := \mathbb{E}[Y | R] - \text{MaxLoss}$

$$\mathbb{E}[\mathbb{I}_{\mathbb{E}[Y|R] > \text{MaxLoss}}]$$

- Digital options $X := S(T) - K$ where S is an asset price satisfying an SDE and K is the strike price

$$\mathbb{E}[\mathbb{I}_{S(T) > K}]$$

- Component failure: $X := g(Y)$ where g depends on the solution of a PDE with random coefficients Y .

$$\mathbb{E}[\mathbb{I}_{g(Y)}]$$

The problem: Computing probabilities

- Financial risk assessment $X := \mathbb{E}[Y | R] - \text{MaxLoss}$

$$\mathbb{E}[\mathbb{I}_{\mathbb{E}[Y|R] > \text{MaxLoss}}] \approx \mathbb{E}\left[\mathbb{I}_{\frac{1}{N} \sum_{i=1}^N Y^{(i)}(R) > \text{MaxLoss}}\right]$$

- Digital options $X := S(T) - K$ where S is an asset price satisfying an SDE and K is the strike price

$$\mathbb{E}[\mathbb{I}_{S(T) > K}] \approx \mathbb{E}[\mathbb{I}_{S_h(T) > K}]$$

where S_h is an Euler-Maruyama or Milstein approximations with step size h .

- Component failure: $X := g(Y)$ where g depends on the solution of a PDE with random coefficients Y .

$$\mathbb{E}[\mathbb{I}_{g(Y)}] \approx \mathbb{E}[\mathbb{I}_{g_h(T)}]$$

where g_h is a Finite Element approximation with grid size h .

Monte Carlo: A General Framework

Focus on

$$\mathbb{E}[f(X)]$$

for some function f . For our setup, $f(X) := \mathbb{I}_{X>0}$. Assume we can approximate $X \approx X_\ell$ with $\ell \in \mathbb{N}$

Assumptions

- Work of X_ℓ is $\propto 2^{\gamma\ell}$.
- Bias: $E_\ell := |\mathbb{E}[f(X_\ell) - f(X)]| \propto 2^{-\alpha\ell}$.

When the dimensionality of X is high, best option is to use Monte Carlo

$$\mathbb{E}[\mathbb{I}_{X>0}] \approx \frac{1}{M} \sum_{m=1}^M \mathbb{I}_{X_L^{(m)}>0}$$

To approximate $\mathbb{P}[X > 0]$ with an error tolerance ε , need $M = \mathcal{O}(\varepsilon^{-2})$ and $L = \mathcal{O}(\frac{1}{\alpha} |\log \varepsilon|)$ hence complexity is $\mathcal{O}(\varepsilon^{-2-\gamma/\alpha})$.

Multilevel Monte Carlo: A General Framework

The MLMC estimator is based on

$$\begin{aligned}\mathbb{E}[f(\mathbf{X})] &= \mathbb{E}[f(\mathbf{X}_0)] + \sum_{\ell=1}^{\infty} \mathbb{E}[f(\mathbf{X}_\ell) - f(\mathbf{X}_{\ell-1})] \\ &\approx \mathbb{E}[f(\mathbf{X}_0)] + \sum_{\ell=1}^L \mathbb{E}[f(\mathbf{X}_\ell) - f(\mathbf{X}_{\ell-1})] \\ &\approx \frac{1}{M_0} \sum_{m=1}^{M_0} f(\mathbf{X}_0^{0,m}) + \sum_{\ell=1}^L \frac{1}{M_\ell} \sum_{m=1}^{M_\ell} f(\mathbf{X}_\ell^{\ell,m}) - f(\mathbf{X}_{\ell-1}^{\ell,m})\end{aligned}$$

Multilevel Monte Carlo: A General Framework

Assumptions

- Work of X_ℓ is $W_\ell \propto 2^{\gamma\ell}$.
- Bias: $|\mathbb{E}[f(X_\ell) - f(X)]| \propto 2^{-\alpha\ell}$.
- Variance: $\text{Var}[X_\ell - X_{\ell-1}] \propto 2^{-\beta\ell}$.

Theorem

For **Lipschitz** f , the overall cost of Multilevel Monte Carlo for computing $\mathbb{E}[f(x)]$ to accuracy ε using optimal L , $\{M_\ell\}_{\ell=0}^L$ is

$$\begin{cases} \varepsilon^{-2} & \beta > \gamma \\ \varepsilon^{-2}(\log \varepsilon)^2 & \beta = \gamma \\ \varepsilon^{-2 - \frac{\gamma - \beta}{\alpha}} & \beta < \gamma \end{cases}$$

Example

For a standard European call option we have $\mathbb{E}[f(X)]$ for $X = S(T) - K$ and $f(X) = \max(X, 0)$. Approximating $S(T)$ by Euler-Maruyama satisfies the previous assumptions with $\alpha = \beta = \gamma = 1$. The complexity is

- $\mathcal{O}(\varepsilon^{-3})$ for Monte Carlo.
- $\mathcal{O}(\varepsilon^{-2}(\log \varepsilon)^2)$ using Multilevel Monte Carlo.

Discontinuous f : Key assumptions

Our quantity of interest is $\mathbb{E}[\mathbb{I}_{X>0}]$ is discontinuous, need a different kind of analysis.

Assumptions

For all $\ell \in \mathbb{N}$ define

$$\delta_\ell := \frac{|X_\ell|}{\sigma_\ell} \geq 0,$$

for some random variable $\sigma_\ell > 0$. For all ℓ :

- 1 There is $\delta > 0$ such that for $x \leq \delta$ we have $\mathbb{P}[\delta_\ell \leq x] \lesssim x$.
- 2 There is $q > 2$ such that

$$\left(\mathbb{E} \left[\left(\frac{|X_\ell - X|}{\sigma_\ell} \right)^q \right] \right)^{1/q} \lesssim 2^{-\beta\ell/2}.$$

Lemma

$$\text{Var}[\mathbb{I}_{X>0} - \mathbb{I}_{X_\ell>0}] \lesssim 2^{-\frac{q}{q+1}\beta/2}$$

Proof. $|X - X_\ell| \approx \mathcal{O}(2^{-\ell\beta/2})$ □

Corollary

Computing $\mathbb{E}[\mathbb{I}_{X>0}]$ to accuracy ε using Multilevel Monte Carlo has cost:

$$\begin{cases} \varepsilon^{-2} & \beta > \frac{q+1}{q} \cdot 2\gamma \\ \varepsilon^{-2}(\log \varepsilon)^2 & \beta = \frac{q+1}{q} \cdot 2\gamma \\ \varepsilon^{-2 - (\gamma - \frac{q}{q+1}\beta/2)/\alpha} & \beta < \frac{q+1}{q} \cdot 2\gamma \end{cases}$$

Previous research

- M. B. Giles, D. J. Higham, and X. Mao. “Analysing multi-level Monte Carlo for options with non-globally Lipschitz payoff”. In: *Finance and Stochastics* 13.3 (2009), pp. 403–413
Original analysis of classical MLMC for discontinuous payoffs in SDE example.
- M. B. Giles, T. Nagapetyan, and K. Ritter. “Multilevel Monte Carlo approximation of distribution functions and densities”. In: *SIAM/ASA journal on Uncertainty Quantification* 3.1 (2015), pp. 267–295
Deals with similar problems in the generality of the current work. Uses different method based on smoothing the discontinuity. Assumes differentiability of PDF and requires further analysis to determine effect of smoothing parameter on bias/variance.
- C. Bayer, C. B. Hammouda, and R. Tempone. “Numerical smoothing and hierarchical approximations for efficient option pricing and density estimation”. In: *arXiv preprint arXiv:2003.05708* (2020)
Same as above. Smooths the discontinuity by integrating using a high order method with respect to one of the dimensions.

Previous research (adaptivity)


- D. Elfverson, F. Hellman, and A. Målqvist. “A multilevel Monte Carlo method for computing failure probabilities”. In: *SIAM/ASA Journal on Uncertainty Quantification* 4.1 (2016), pp. 312–330
Selective refinement of samples. Based on relaxing the condition. Assumes uniform almost sure error bounds (works well for PDEs with random coefficients but not stochastic models).
- M. Broadie, Y. Du, and C. C. Moallemi. “Efficient risk estimation via nested sequential simulation”. In: *Management Science* 57.6 (2011), pp. 1172–1194
Adaptive sampling for nested expectation with Monte Carlo methods.
- M. B. Giles and A.-L. Haji-Ali. “Multilevel nested simulation for efficient risk estimation”. In: *SIAM/ASA Journal on Uncertainty Quantification* 7.2 (2019), pp. 497–525. DOI: [10.1137/18M1173186](https://doi.org/10.1137/18M1173186)
Adaptive sampling for MLMC applied to nested expectations only. Requires stronger conditions on the random variables than here.

Adaptive Multilevel Monte Carlo: Algorithm

Refine samples of X_ℓ to $X_{\ell+\eta_\ell}$, where $0 \leq \eta_\ell \leq \lceil \theta \ell \rceil$ is the smallest integer for which

$$\delta_{\ell+\eta_\ell} \geq 2^{\frac{\gamma}{r}(\theta \ell(1-r) - \eta_\ell)}$$

for constants $r > 1$ and $0 \leq \theta \leq 1$. Recall that $\delta_\ell := \frac{|X_\ell|}{\sigma_\ell} \geq 0$.




Adaptive Multilevel Monte Carlo: Algorithm

Refine samples of X_ℓ to $X_{\ell+\eta_\ell}$, where $0 \leq \eta_\ell \leq \lceil \theta \ell \rceil$ is the smallest integer for which

$$\delta_{\ell+\eta_\ell} \geq 2^{\frac{\gamma}{r}(\theta \ell(1-r) - \eta_\ell)}$$

for constants $r > 1$ and $0 \leq \theta \leq 1$. Recall that $\delta_\ell := \frac{|X_\ell|}{\sigma_\ell} \geq 0$.




Adaptive Multilevel Monte Carlo: Algorithm

Refine samples of X_ℓ to $X_{\ell+\eta_\ell}$, where $0 \leq \eta_\ell \leq \lceil \theta \ell \rceil$ is the smallest integer for which

$$\delta_{\ell+\eta_\ell} \geq 2^{\frac{\gamma}{r}(\theta \ell(1-r) - \eta_\ell)}$$

for constants $r > 1$ and $0 \leq \theta \leq 1$. Recall that $\delta_\ell := \frac{|X_\ell|}{\sigma_\ell} \geq 0$.



Adaptive Multilevel Monte Carlo: Analysis

Theorem

There is $\bar{r} > 1$ such that for $1 < r < \bar{r}$:

- The expected work of sampling $\mathbb{I}_{X_{\ell+\eta_\ell} > 0}$ is $W_\ell \propto 2^{\gamma\ell}$.
- The variance is

$$\text{Var}[\mathbb{I}_{X_\ell > 0} - \mathbb{I}_{X_{\ell+\eta_\ell} > 0}] \propto 2^{-\frac{q}{q+1} \frac{1+\theta}{2} \beta \ell}$$

for

$$\theta = \begin{cases} \frac{1}{2^{\frac{q+1}{q} \frac{\gamma}{\beta}} - 1} & \beta < \frac{q+1}{q} \gamma \\ 1 & \beta > \frac{q+1}{q} \gamma \end{cases}.$$

Adaptive Multilevel Monte Carlo: Complexity

Corollary

Computing $\mathbb{E}[\mathbb{I}_{X>0}]$ to accuracy ε using (non-)adaptive Multilevel Monte Carlo has cost:


Non-Adaptive:

$$\begin{cases} \varepsilon^{-2} & \beta > \frac{q+1}{q} \cdot 2\gamma \\ \varepsilon^{-2}(\log \varepsilon)^2 & \beta = \frac{q+1}{q} \cdot 2\gamma \\ \varepsilon^{-2 - (\gamma - \frac{q}{q+1}\beta/2)/\alpha} & \beta < \frac{q+1}{q} \cdot 2\gamma \end{cases}$$

Adaptive:

$$\begin{cases} \varepsilon^{-2} & \beta > \frac{q+1}{q} \cdot \gamma \\ \varepsilon^{-2}(\log \varepsilon)^2 & \beta = \frac{q+1}{q} \cdot \gamma \\ \varepsilon^{-2 - (\gamma - \frac{q}{q+1}\beta(1+\theta)/2)/\alpha} & \beta < \frac{q+1}{q} \cdot \gamma \end{cases}$$

Work/variance proof idea



Example

When approximating the price of a digital option

$$\mathbb{E}[\mathbb{I}_{S(T) > K}],$$


using Euler-Maruyama approximation of $S(T)$, or the risk estimation problem

$$\mathbb{E}[\mathbb{I}_{\mathbb{E}[Y|R] > 0}],$$


the assumptions hold for $\alpha = \beta = \gamma = 1$ and any $q < \infty$. The complexity is (for any $\nu > 0$)

- $\mathcal{O}(\varepsilon^{-3})$ for Monte Carlo.
- $o(\varepsilon^{-2.5-\nu})$ for non-adaptive Multilevel Monte Carlo.
- $o(\varepsilon^{-2-\nu})$ for adaptive Multilevel Monte Carlo.


Numerical Test: Digital Options



Numerical Test: Digital Options



Numerical Test: Digital Options



Other points in the paper

In <https://arxiv.org/abs/2107.09148>

- Nested expectation: Differs from previous work in that the same samples are used for computing the refinement, η_ℓ and for computing the estimate $X_{\ell+\eta_\ell}$. Leads to reduced cost and more relaxed assumptions.
- Discussion on choices of σ_ℓ in nested expectation.
- Motivation of previous assumptions in the case of nested expectation and SDEs.
- Analysis of weak error bounds and corresponding necessary assumptions.

Conclusion

- Accurate computation of probabilities by standard Monte Carlo techniques is expensive when the underlying observable must be approximated for each sample.
- Multilevel Monte Carlo is a great method to reduce this cost, but suffers for probabilities due to the intrinsic discontinuity.
- Adaptive sampling provides a general framework to improve Multilevel Monte Carlo performance for probabilities, in many cases to optimal $\mathcal{O}(\varepsilon^{-2})$ cost.
- Other applications: Barrier options and computing sensitives.
- See also special session: “Monte Carlo Methods for Discontinuous Functions”, Wednesday 9am to 11am (UTC+2).

Numerical Tests: Digital Options

For constant $\mu, \sigma, S(0)$ consider the asset

$$dS(t) = \mu S(t) dt + \sigma S(t) dW(t).$$

Compute

$$\mathbb{E}[\mathbb{I}_{X>0}] := \mathbb{E}[\mathbb{I}_{S(T)-K>0}]$$

for some strike price $K > 0$. We use Euler-Maruyama with a step size $h_\ell = 2^{-\ell}$ to approximate $S_{h_\ell}(\cdot) \approx S(\cdot)$ and set

$$X_\ell := S_{h_\ell}(T) - K.$$

The assumptions are satisfied using constant $\sigma_\ell \equiv 1$ for $\alpha = \beta = \gamma = 1$ and any $q < \infty$ giving complexity $\mathcal{O}(\varepsilon^{-2.5-\nu})$ for standard Multilevel Monte Carlo and $\mathcal{O}(\varepsilon^{-2-\nu})$ for any $\nu > 0$ using adaptive Multilevel Monte Carlo.

Numerical Tests: Digital Options

Consider the assets

$$dS^{(i)}(t) = \mu^i S^{(i)}(t) + \sigma^i S^{(i)}(t) dW^{(i)}(t)$$

where

$$W^i(t) = \rho W_{\text{com}}^{(i)}(t) + \sqrt{1 - \rho^2} W_{\text{ind}}^{(i)}(t)$$

for $1 \leq i \leq 10$. Consider the digital option with payoff

$$\mathbb{I}_{\left(\frac{1}{10} \sum_{i=1}^{10} S^{(i)}(t)\right) > K}.$$

Thus, compute

$$\mathbb{E} \left[\mathbb{I}_{\left(\frac{1}{10} \sum_{i=1}^{10} S^{(i)}(t)\right) > K} \right].$$